


JPL: Duane Waliser, Yi Chao, Qinbin Li, Annmarie Eldering UCLA: Jinwon Kim, Kuo-Nan Liou, Yongkang Xue, Alex Hall, Mimi Hughes, Sarah Kapnick, Robert Fovell, Jim McWilliams

5th Annual California Climate Change Conference Sacramento Convention Center September 8 - 10, 2008





- UCLA + JPL Joint Institute
- JIFRESSE Research Foci & Objectives
- Regional Model Development & Framework

# Outline

- Today's California Climate Change Themes
  - Hydroclimate (Kim et al. 2008; POSTER 4)
  - □ Sierra snowpack & snow physics (Kapnick and Hall, 2008; Waliser et al. 2008; POSTERS 54 & 5x?)
  - □ Santa Ana wind conditions (Hughes et al. 2008; POSTER 45)



# California Climate Change: UCLA+JPL

In 2006, UCLA & JPL established a <u>Joint Institute for</u>
Regional Earth System Science & Engineering (JIFRESSE)
to promote, stimulate, and engage UCLA (*Modeling*) and
JPL (*Observations*) in cutting edge Earth System science
research to:




# California Climate Change: UCLA+JPL


In 2006, UCLA & JPL established a <u>Joint Institute for</u>
Regional Earth System Science & Engineering (JIFRESSE)
to promote, stimulate, and engage UCLA (*Modeling*) and
JPL (*Observations*) in cutting edge Earth System science
research to:

- Increase understanding about factors that affect climate-related environmental changes, with particular attention to regional/CA issues;
- Support the design of future JPL/NASA space missions & observation networks related to detecting, monitoring, and projecting climate changes;
- Enhance the educational mission of UCLA through collaboration with JPL.

# California Climate Change: Motivation


IPCC AR4 model projections agree that California will warm in this century but disagree on whether it will become wetter/drier. This implies that some physical processes are inadequately represented in GCMs.






Apply our unique strengths in system engineering and observations (JPL) and process understanding and modeling (ULCA) to improve our capabilities to detect and predict changes in California's climate and ecosystems and contribute to the State's awareness and understanding, and adaptation and mitigation strategies.

Kapnick & Hall, 2008



✓ A snow water equivalent dataset was developed from 1930 -2007.

Kapnick & Hall, 2008




- ✓ A snow water equivalent dataset was developed from 1930 -2007.
- ✓ For each winter season, the date of peak snow mass was calculated.

Kapnick & Hall, 2008




- ✓ For each winter season, the date of peak snow mass was calculated.
- ✓ Trend: date of peak snow mass occurs earlier by 0.4 days per decade.



Kapnick & Hall, 2008



- ✓ For each winter season, the date of peak snow mass was calculated.
- ✓ Trend: date of peak snow mass occurs earlier by 0.4 days per decade.




Based on these results and considerations of temperature projections, the date of peak snow mass would be expected to occur 3-9 days earlier by end of century.

### California Snowpack Projections: Model Uncertainties

What do the models (directly) project?

#### **IPCC AR4 Projections**




**But how realistic?** 


### California Snowpack Projections: Model Uncertainties

What do the models (directly) project?

#### **IPCC AR4 Projections**




**But how realistic?** 




### California Snowpack Projections: Model Uncertainties

What do the models (directly) project?

#### **IPCC AR4 Projections**



**But how realistic?** 





RESMs – and their dynamic downscaling of GCMs - afford:


- □ greater spatial resolution [O(1-10km)]
- inclusion of more processes and interactions
- closer connections to societal impacts
- □ a complementary role to GCMs
- a numerical laboratory for GCM parameterization development

RESMs – and their dynamic downscaling of GCMs - afford:


- □ greater spatial resolution [O(1-10km)]
- inclusion of more processes and interactions
- closer connections to societal impacts
- □ a complementary role to GCMs
- a numerical laboratory for GCM parameterization development








Ongoing Atmos-Ocean & Ocean-Bio Coupling Studies & Forecasts



Ongoing
Hydrology and Synotpic
Studies e.g. Today's Snowpack
& Santa Ana Wind
Studies




Climate Change Experiments

#### **Forcing**


Climate Change
Projection
NCAR CCSM3 –
SRESA1B
Scenario
1971-1980
&
2045-2054




Climate Change Experiments




Climate Change Experiments



Hydroclimate Changes: Oct-March; ~2050 vs 1975; 36km




Hydroclimate Changes: Oct-March; ~2050 vs 1975; 36km




**Warming**: 0° to 2.5°C. Larger with altitude.

Hydroclimate Changes: Oct-March; ~2050 vs 1975; 36km



Hydroclimate Changes: Oct-March; ~2050 vs 1975; 36km




**Warming**: 0° to 2.5°C. Larger with altitude.

**SWE:** Up to -80%

Albedo Decreases: Up to -50%. Implies a snow-albedo impact on temperature signal.

Hydroclimate Changes: Oct-March; ~2050 vs 1975; 36km



**Warming**: 0° to 2.5°C. Larger with altitude.


**SWE:** Up to -80%

Albedo Decreases: Up to -50%. Implies a snow-albedo impact on temperature signal.

Runoff: Up to -80%.

Negatively impacting water resouces in California

Hydroclimate Changes: Oct-March; ~2050 vs 1975; 36km



**Warming**: 0° to 2.5°C. Larger with altitude.

**SWE:** Up to -80%


Albedo Decreases: Up to -50%. Implies a snow-albedo impact on temperature signal.


Runoff: Up to -80%.

Negatively impacting water resouces in California

Snow physics modeling is important


Impact of Model Resolution/Topography






**LOW ELEVATION** 

Impact of Model Resolution/Topography







- SWE is generally less in projected future climate.
- Differences in 12km vs 36km are not due to differences in changes to snowfall.

**LOW ELEVATION** 

Impact of Model Resolution/Topography







Greater reduction in snowpack in 36km => likely due to greater snow -albedo feedback since overall less snow.

**LOW ELEVATION** 

Impact of Model Resolution/Topography







12km Resolution:
Snowpack loss is greater
at lower elevations =>
likely due to greater snow
-albedo feedback as snow
is less in lower elevations

**LOW ELEVATION** 

Impact of Model Resolution/Topography







Assessements of SWE need to be derived from high -resolution model projections.

**LOW ELEVATION** 


Impact of Snow Albedo/Aerosol Deposition: Role of Emissions?




#### SWE sensitivity to changes in snow albedo



Impact of Snow Albedo/Aerosol Deposition: Role of Emissions?




#### SWE sensitivity to changes in snow albedo




Modest increases in snow albedo lead to significant inceases in SWE later in season

Impact of Snow Albedo/Aerosol Deposition: Role of Emissions?




#### SWE sensitivity to changes in snow albedo




Modest decreases in snow albedo lead to significant decreases in SWE earlier in season

Impact of Snow Albedo/Aerosol Deposition: Role of Emissions?



#### SWE sensitivity to changes in snow albedo



Small changes in snow albedo have dramatic changes on projected SWE.


Needs to be examined more thoroughly with respect to projected aerosol emission.

Single versus multi-layer snow model

**Forcing** 


Sensitivity
Simulation
NCEP/NCAR
Reanalysis
1998


Single versus multi-layer snow model




# California Climate Change: Dynamical Downscaling Single versus multi-layer snow model **Forcing Model Results Sensitivity Simulation Modeled Snow Layers** NCEP/NCAR 1-layer Multi-layer Reanalysis **80km Resolution** 1998


Impact of Snow Layer Physics Representation






Impact of Snow Layer Physics Representation






Significant Reductions in modeled SWE biases – typically an over estimate.


Impact of Snow Layer Physics Representation






Significant Reductions in modeled SWE biases – typically an over estimate.

Impact of Snow Layer Physics Representation






Significant Reductions in modeled SWE biases – typically an over estimate.

Less realistic single layer snow models may underestimate projected snowpack loss.


#### California Climate Change: Santa Ana Winds Hughes et al. 2008





#### California Climate Change: Santa Ana Winds Hughes et al. 2008





- Analysis of a regional climate hindcast'indicates Santa Ana wind conditions have decreased 30-50% from 1959 to 2001.
- Reduction arises from both large-scale synoptic forcing (strong offshore upper level winds) and local katabatic forcing (cold desert air pouring through the gaps).
- The climate change dynamic downscaling experiment suggests further decreases in the future (~2050).

# Summary

- An trend in earlier Sierra snowmelt timing is present in monthly SWE observations attributed to the sensitivity of peak snow mass date to local March temperature.
- Regional dynamic downscaling experiments have been performed based on NCAR CCSM SRESA1B suggesting considerable changes to CA hydroclimate by 2050.
- There is considerable model sensitivity of CA snowmelt and snowpack loss to model resolution, snow albedo treatment, and model snow layer formulation.
- Santa Ana winds are less frequent under climate warming conditions because the continental interior warms more than ocean, altering large-scale and local pressure gradients.
- □ Caveat: Some results based on only one GCM projection & one RESM formulation.

# Future Work

- Regional model validation with ground-based and satellite observations where possible: synoptic fields (e.g., temperature, clouds, winds), hydroclimate (e.g. snowcover, albedo), air quality.
- Continue work on improving snow physics, e.g., multi-layer formulation, snow spectral albedo parameterization based on snow grain size and black carbon/dust contamination.
- Compare observational dataset of peak snow mass date to regional model output both for validation purposes and for understanding mechanisms for the changes.
- □ Santa Ana Winds: 1) Validate analysis against ground-based observations, and 2) Identify how other critical fire weather parameters (e.g., relative humidity) have changed.
- Investigation of the impact of anthropogenic climate change on the air quality in California.

  www.jifresse.ucla.edu